

Fresh Tracheostomy Care Guideline

Inclusion Criteria:

- All postoperative fresh tracheostomy patients
- Any previous tracheostomy where the stoma has been surgically manipulated
- All transfers of fresh tracheostomy patients performed in the past 10 days

Exclusion Criteria:

- Simple tracheostomy revision

To PICU/ CVICU/
NICU postop

Assessment/Diagnostics

- Vital signs per standard of care
- Cardiorespiratory monitoring – continuous
- Portable CXR per order
- Document trach tube size & type
- If abnormal anatomy or trach position, anatomic diagram **MUST be placed** at bedside
- Stay sutures must be clearly marked right and left and taped on respective sides of chest for easy access.
- Assess skin under and around trach & trach ties with each routine assessment

Interventions/Treatment

- IV fluids initially; advance feeding as clinically indicated
Cefazolin 25 mg/kg/dose IV q 8hr (<40 kg);
1000 mg IV q 8 hr (>40 kg) (Max: 6 gm/day) Duration:
up to 24 hours **OR**
Clindamycin 10 mg/kg/dose IV q6hr (<60kg);
600 mg IV q 6hr (>60kg) (Max: 4.8 gm/day) Duration: up to 24
hours **AND** gentamicin 2.5 mg/kg IV q8h < 40 kg, 100 mg IV
q8h (40-60 kg) and 120 mg IV q8h > 60 kg
- Oxygen via trach collar to keep sats appropriate for baseline
- Ventilator management by ICU service including non-invasive CO₂ measurement monitoring when indicated
- Humidification via trach collar
- Notify Trach Specialty Nurse
- Tracheostomy Care per CHOC Policy: Tracheostomy tube: stoma care and tie change.
- Extra trach at bedside (same size & one smaller) and including transfers/transports, additional equipment as specified in CHOC Policy
- Hyperextension of neck for appropriate patients
- First trach change by ENT surgeon before transfer out of ICU

Continued Considerations

- Obtain Pulmonary consult prior to discharge

At High Risk For:

- Ripping of Stay Sutures
- Trach/airway plugging
- Device-related pressure injuries

Recommendations/ Considerations

- Common indications for tracheostomy are: prolonged intubation, pulmonary toilet, upper airway obstruction, craniofacial syndromes, neurologic impairment, trauma, vocal cord paralysis
- Early complications may include: bleeding, tube displacement, tube blockage, subcutaneous & mediastinal emphysema, pneumothorax, and rarely esophageal perforation & nerve damage.
- Antibiotics for surgical infection prophylaxis should be continued up to 24 hrs.
- Pediatric tracheostomies are typically changed between postop day 5 to 10.
- Stay sutures are generally removed between postop day 5 to 10.
- There must be ready access to tracheostomy and intubation trays in the ICUs
- Trach patients are at high risk for device-related pressure ulcers and skin breakdown. Care planning and interventions should be implemented as appropriate.
- Use Mepilex transfer dressing around trach

Patient Education

- Trach care
- Suctioning technique
- CPR Education
- Tracheostomy Home Care Instructions located on PAWS

References

Fresh Tracheostomy Care Guideline

Carron JD, Derkay CS, et al. Pediatric Tracheotomies: Changing Indications and Outcomes. *The Laryngoscope* 110: July 2000, pp 1099-1104.
<http://www.laryngoscope.com/pt/re/laryngoscope/abstract.00005537-200007000-00006.htm;jsessionid=JBvYvSgWgBTnfnzLGXpZv3d2yRzyYmMsGb1zSTQbSnxxKf5nnyVq!-26702612!181195628!8091!-1>

Darden D, Towbin R, et al. Pediatric Tracheotomy: Is Postoperative Chest X-Ray Necessary? *Annals of Otolaryngology, Rhinology & Laryngology* 110: April 2001, pp 345-348.
http://www.annals.com/toc/auto_abstract.php?id=13055

Gluth MB, Maska S, et al. Postoperative Management of Pediatric Tracheostomy: Results of a Nationwide Survey. *Otolaryngology – Head and Neck Surgery* 122: May 2000, pp 701-705. http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WP4-4HGNP06-H&_user=10&_coverDate=05%2F31%2F2000&_alid=839433512&_rdoc=2&_fmt=high&_orig=search&_cdi=6980&_sort=d&_docanchor=&_view=c&_ct=2&_acct=C000050221&_version=1&_urVersion=0&_userid=10&md5=b33c31e65d77b005f8896225dd192e58

Mitchell RB, Hussey HM. Clinical Consensus Statement: Tracheostomy Care. *American Academy of Otolaryngology – Head and Neck Surgery*. September, 2012.
<http://oto.sagepub.com/content/early/2012/09/18/0194599812460376>

Wootten CT, French LC, et al. Tracheotomy in the First Year of Life: Outcomes in Term Infants, the Vanderbilt Experience. *Otolaryngology – Head and Neck Surgery* 134: March 2006, pp 365-369. http://www.sciencedirect.com/science?_ob=ArticleURL&_udi=B6WP4-4J9WPS6-B&_user=10&_coverDate=03%2F31%2F2006&_alid=839433616&_rdoc=8&_fmt=high&_orig=search&_cdi=6980&_sort=d&_docanchor=&_view=c&_ct=12&_acct=C000050221&_version=1&_urVersion=0&_userid=10&md5=d7b3fe9befa0d93eb21c878c06d9ed42

CHOC Policy

CHOC Tracheostomy Home Care Instructions, February, 2010.