

# Pyelonephritis Care Guideline

**Inclusion Criteria:** Previously healthy children 1 month to 5 years of age with an abnormal UA suspicious for infection who:

- require IV fluids
- require IV antibiotics due to severe disease
- are intolerant of oral antibiotics, are failing oral antibiotic treatment
- are not deemed eligible for outpt management (social situation, uncertainty of diagnosis, etc)

**Exclusion Criteria:** Children < 1 month and > 5 years of age, PICU status, known major genitourinary anomalies, significant co-morbid condition(s) (e.g. sepsis, meningitis, spina bifida)

**Assessment:** Hydration status, signs of sepsis, CVAT, flank tenderness, urinalysis, urine c&s, blood culture if indicated

**Empiric antibiotics:**  
Ceftriaxone 75mg/kg/day IV once daily (max 2 gm)  
Adjust per culture results and response to therapy

**Imaging Studies** (if not previously done)  
Renal ultrasound while inpatient  
VCUG generally in 2-3 weeks as outpatient if renal ultrasound suggests high grade VUR or obstructive uropathy

**Discharge Criteria**

- Clinically improving
- Able to tolerate po
- Outpatient follow up in place
- Families and clinicians to have high index of suspicion and consider recurrent infection with subsequent febrile illness

## Recommendations/Considerations

- Timely diagnosis & treatment of urinary tract infections results in a lower incidence of renal scarring, hypertension, & loss of renal function.
- Pyelonephritis can be generally presumed with the presence of fever and a suspicious urinalysis (+/- the presence of upper tract signs such as vomiting, flank pain, costovertebral angle tenderness etc.).
- A well done prospective randomized controlled multi-center study on 1-24 month old children demonstrates that pyelonephritis, in general, can be managed on an outpt basis (Hoberman, et al. See references).
- Sensitivity & specificity of both the UA and culture are dependent on the method of collection & evaluation.
- The distinction between simple urinary tract infection & pyelonephritis can impact the need for imaging in girls.
- A normal late prenatal ultrasound (after 30-32 weeks gestation) done in an experienced center may preclude the need for a renal ultrasound.
- Consider inpatient VCUG for recurrent febrile UTI, concern for posterior urethral valves or non-compliant parent/family/ social situation .
- Antibiotics should be continued for at least 10 days.
- The use of antibiotic prophylaxis is considered controversial in the literature.

## Patient Education

KidsHealth handout for Urinary Tract Infection – Parent Version

## References

### Pyelonephritis Care Guideline

American Academy of Pediatrics, Subcommittee on Urinary Tract Infection. Clinical Practice Guideline for the Diagnosis and Management of the Initial UTI in Febrile Infants and Children 2-24 months. *Pediatrics*. September 2011, 128(3) 595-610.  
<http://pediatrics.aappublications.org/cgi/content/abstract/128/3/595>

Bauer R, Kogan B. New Developments in the Diagnosis and Management of Pediatric UTIs. *Urologic Clinics of North America*, 2008 (35): 47-58.  
[http://www.urologic.theclinics.com/article/S0094-0143\(07\)00092-4/abstract](http://www.urologic.theclinics.com/article/S0094-0143(07)00092-4/abstract)

Hoberman A, et al. Oral Versus Initial Intravenous Therapy for Urinary Tract Infections in Young Febrile Children. *Pediatrics*, 104(1): 79-86, 1999.  
<http://pediatrics.aappublications.org/cgi/content/abstract/104/1/79>

Hoberman A, et al. Imaging Studies after a First Febrile Urinary Tract Infection in Young Children. *New England Journal of Medicine*. January 2003, 348(3) 195-202.  
<http://content.nejm.org/cgi/content/abstract/348/3/195>

Larcombe J. Urinary Tract Infection in Children. *BMJ Clinical Evidence* [Online] December 2006. <http://www.clinicalevidence.bmj.com/ceweb/conditions/chd/0306/0306-get.pdf>.

Montini G, et al. Febrile Urinary Tract Infections in Children. *New England Journal of Medicine*. July 2011; 365:239-250.  
<http://www.nejm.org/doi/full/10.1056/NEJMra1007755>

Raszka W, Khan O. Pyelonephritis. *Pediatrics in Review*. October 2005, 26(10) 364-370.  
<http://pedsinreview.aappublications.org/cgi/content/full/26/10/364>

Shaikh N, Hoberman A. Acute Management, Imaging, and Prognosis of Urinary Tract Infections in Children. *UpToDate* [Online] February 2008.

UTI Guideline Team, Cincinnati Children's Hospital Medical Center: Evidence-based care guideline for medical management of first urinary tract infection in children 12 years of age or less, Guideline 7, pages 1-23, November 2006.  
<http://www.cincinnatichildrens.org/svc/alpha/h/health-policy/ev-based/uti.htm>