

Junctional Ectopic Tachycardia (JET) Care Guideline for Cardiovascular Intensive Care Unit (CVICU)



Inclusion Criteria: Postoperative Cardiac Surgical Patient. Diagnosis of JET confirmed by EKG. Patient must have hemodynamic instability due to the arrhythmia.

Exclusion Criteria: Patients without temporary pacing wires

Initial Management

- Adequate sedation
- Electrolytes (especially magnesium)
- Temperature 36°C - 37°C rectal
- Decrease vasoactives (especially dopamine)
- Attempt brief period of atrial pacing over ventricular rate
- Check threshold of wires
- Obtain atrial ECG to document rhythm
- Obtain ECHO to rule out effusion and check function
- Notify Cardiologist

Recommendations/ Considerations

The **GOAL** of treatment is to decrease JET rate just low enough to pace over

- Lower medication doses may be appropriate in preterm neonate or impaired renal function

Secondary Medications/Interventions

Frequently reassess rhythm to determine ongoing need for medication

Consider cooling patient further down to 35°C - 36°C rectal

- **Dexmedetomidine** infusion start at 0.2 mcg/kg/hr up to max of 1 mcg/kg/hr.
 - Reserve dexmedetomidine load for extreme situations
- **Amiodarone** 1 mg/kg over 10 minutes. Repeat in 1mg/kg increments to a total dose of 5 mg/kg.
 - Then amiodarone infusion at 5 mcg/kg/min.
 - Give intravenous CaCl prior to amiodarone load
- **Procainamide** 7-10 mg/kg load over 30 minutes
 - Then procainamide infusion at 40-60 mcg/kg/min
 - Check levels at 4 hours
 - Use with caution if ejection fraction already decreased
- **Esmolol** infusion at 50-300 mcg/kg/min
 - Consider loading with caution
 - Consider esmolol as first line agent in patients with JET presenting on POD 2 or greater and the patient is not on dopamine and ejection fraction is normal

Consider EP consult if patient does not respond to first pharmacologic agent

Tertiary Medications/Interventions

- **Sotalol**
- **Caution:** Must have an EP specialist consultation prior to initiation of Sotalol
- Strong caution if patient has received Amiodarone
- Give 1mg/kg intravenous Sotalol over 1 hour. Can consider a second dose in consultation with EP specialist

References for Junctional Ectopic Tachycardia (JET) Care Guideline

- Andreasen JB, Johnson SP, Ravn HB. Junctional ectopic tachycardia after surgery for congenital heart disease in children. *Intensive Care Med.* 2008;34(5):895-902. doi:10.1007/s00134-007-0987-2.
- Batra AS, Chun DS, Johnson TR, et al. A prospective analysis of the incidence and risk factors associated with junctional ectopic tachycardia following surgery for congenital heart disease. *Pediatr Cardiol.* 2006;27(1):51-55. doi:10.1007/s00246-005-0992-6.
- Chrysostomou, C, Beerman L, Shiderly D, Berry D, Morell VO, Munoz R. Dexmedetomidine: a novel drug for the treatment of atrial and junctional tachyarrhythmias during the perioperative period for congenital cardiac surgery: a preliminary study. *Anesth Analg.* 2008;107(5):1514-1522. doi:10.1213/ane.0b013e318186499c.
- Dodge-Khatami A, Miller OI, Anderson RH, Gil-Jaurena JM, Goldman AP, de Leval MR. Impact of junctional ectopic tachycardia on postoperative morbidity following repair of congenital heart defects. *Eur J Cardiothorac Surg.* 2002;21(2):255-259. doi:10.1016/S1010-7940(01)01089-2.
- Dodge-Khatami, A, Miller OI, Anderson, RH, et al. Surgical substrates of postoperative junctional ectopic tachycardia in congenital heart defects. *J Thorac Cardiovasc Surg.* 2002;123(4):624-630. doi:10.1067/mtc.2002.121046.
- Dorman BH, Sade RM, Burnette JS, et al. Magnesium supplementation in the prevention of arrhythmias in pediatric patients undergoing surgery for congenital heart defects. *American Heart Journal.* 2000;139(3):522-528. doi:10.1016/S0002-8703(00)90097-8.
- Grosse-Wortmann L, Kreitz S, Grabitz RG, et al. Prevalence of and risk factors for perioperative arrhythmias in neonates and children after cardiopulmonary bypass: continuous holter monitoring before and for three days after surgery. *J Cardiothorac Surg.* 2010;5:85. doi:10.1186/1749-8090-5-85.
- Hoffman TM, Bush DM, Wernovsky G, et al. Postoperative junctional ectopic tachycardia in children: incidence, risk factors, and treatment. *Ann Thorac Surg.* 2002;74(5):1607-1611. doi:10.1016/S0003-4975(02)04014-6.
- Kendrick JG, Macready JJ, Kissoon N. Amiodarone treatment of junctional ectopic tachycardia in a neonate receiving extracorporeal membrane oxygenation. *Ann Pharmacother.* 2006;40(10):1872-1875. doi:10.1345/aph.1H148.
- Kovacikova L, Hakacova N, Dobos D, Skrak P, Zahorec M. Amiodarone as a first-line therapy for postoperative junctional ectopic tachycardia. *Ann Thorac Surg.* 2009;88(2):616-622. doi:10.1016/j.athoracsur.2009.04.088.
- Manrique AM, Arroyo M, Lin Y, et al. Magnesium supplementation during cardiopulmonary bypass to prevent junctional ectopic tachycardia after pediatric cardiac surgery: a randomized controlled study. *J Thorac Cardiovasc Surg.* 2010;139(1):162-169. doi:10.1016/j.jtcvs.2009.07.064.
- Mildh L, Hiippala A, Rautiainen P, Pettila V, Sairanen H, Happonen J. Junctional ectopic tachycardia after surgery for congenital heart disease: incidence, risk factors and outcome. *Eur J Cardiothorac Surg.* 2011;39(1):75-80. doi:10.1016/j.ejcts.2010.04.002
- Perry JC, Fenrich AL, Hulse JE, Trienman JK, Friedman RA, Lamberti JJ. Pediatric use of intravenous amiodarone: efficacy and safety in critically ill patients from a multicenter protocol. *J Am Coll Cardiol.* 1996;27(5):1246-1250. doi:10.1016/0735-1097(95)00591-9.
- Valdes SO, Landstrom AP, Schneider AE, Miyake CY, de la Uz CM, Kim JJ. Intravenous sotalol for the management of postoperative junctional ectopic tachycardia. *HeartRhythm Case Rep.* 2018;4(8):375-377. doi:10.1016/j.hrcr.2018.05.007.
- Saul JP, Scott WA, Brown S, et al. Intravenous amiodarone for incessant tachyarrhythmias in children: a randomized, double-blind anti-arrhythmic drug trial. *Circulation.* 2005;112(22):3470-3477. doi:10.1161/CIRCULATIONAHA.105.534149.
- Walsh EP, Saul JP, Sholler GF, et al. Evaluation of a staged treatment protocol for rapid automatic junctional tachycardia after operation for congenital heart disease. *J Am Coll Cardiol.* 1997;29(5):1046-1053. doi:10.1016/S0735-1097(97)00040-5.