


Inclusion criteria: children  $\geq$  3 yrs with headaches

Is urgent emergency department, neuroimaging, or Neurology consultation indicated?



**Recommendations for All Headaches**

**Headache hygiene**

- Regular bedtime and awakening time
- Regular daily exercise
- Good hydration
- Avoid known triggers

**Non-medical interventions**

- Ice pack
- Warm bath
- Nap in a cool, dark room
- Neck and back massage
- Take a walk

- If fails headache hygiene and at least one preventative treatment (min. 4 week trial), refer to Neurology.
- If responds to preventative treatment, continue for at least 3 months and re-evaluate

Characterize Headache Types

See page 2

## Characterizing Headache Types

### Migraine Headache

1. At least 5 attacks fulfilling features # 2-4
2. Headache attack lasting 1 to 72 hrs
3. Headache has at least 2 of the following features:
  - either bilateral or unilateral (frontal/temporal) location
  - pulsating quality
  - moderate to severe intensity
  - aggravated by routine physical activities
4. At least 1 of the following accompanies headache:
  - nausea and/or vomiting
  - photophobia and/or phonophobia (may be inferred from their behavior)

See page 3 for interventions

### Tension Headache

1. Headache lasting from 30 mins to 7 days
2. Headache has at least 2 of the following characteristics:
  - bilateral location
  - pressing/tightening (non-pulsating quality)
  - mild or moderate intensity
  - not aggravated by routine physical activity such as walking or climbing stairs
3. Both of the following:
  - no nausea or vomiting (anorexia may occur)
  - no more than one of photophobia or phonophobia

See page 4 for interventions

### Cluster Headache

1. At least 5 attacks fulfilling criteria # 2-4
2. Severe or very severe unilateral orbital, supraorbital and/or temporal pain lasting 15-180 minutes if untreated
3. Headache is accompanied by at least one of the following:
  - Ipsilateral conjunctival injection and/or lacrimation
  - Ipsilateral nasal congestion and/or rhinorrhea
  - Ipsilateral eyelid edema
  - Ipsilateral forehead and facial sweating
  - Ipsilateral miosis and/or ptosis
  - A sense of restlessness or agitation

See page 4 for interventions

### Analgesic Medication Overuse

1. Headache present on  $\geq 15$  days/month
2. Use of one or more drugs that can be taken for acute and/or symptomatic treatment of headache,  $>3$  times per week over a 3 month period
3. Headache has developed or markedly worsened during medication overuse
4. Headache resolves or reverts to its previous pattern within 2 months after discontinuation of overused medication

See page 4 for interventions

### Sleep Apnea Related Headache

1. Typically a morning headache, bilateral, lasting  $< 4$  hrs and not accompanied with nausea, photophobia, or phonophobia

See page 4 for interventions

# Headache Treatment by Type

## Migraine Headache

### Abortive

#### Common OTC Analgesics

- Indicated for mild-moderate headache.
- NSAIDs tend to be the most effective. NSAIDs dosed at 10-15 mg/kg/dose and acetaminophen at 15-20 mg/kg/dose.
- Should not be used more than twice a week to prevent possible medication overuse headache and toxicity from the analgesic.

#### Sumatriptans

- Indicated for moderate-severe headache. Triptans are **contraindicated** in patients with cardiovascular disease, uncontrolled HTN, basilar migraine, and hemiplegic migraine.
- SUMAtriptan single dose for 6-11 years old is 25 mg. Single dose for  $\geq 12$  years old is 50 mg; can increase to 100 mg if needed. Max dose is 200 mg/24 hours.
- Typically third party payers require SUMAtriptan prior to using any other triptans
- Nasal SUMAtriptan can be used if patient has extreme nausea at onset of headache and inability to tolerate PO. Also useful if not responding well to SUMAtriptan tablets. Dosing guideline 6-12 years old 5-10 mg/dose;  $>12$  years old 10-20 mg/dose. No repeat dose.

#### Other Triptans

- Rizatriptan dosing is 5 mg  $< 40$ kg and 10 mg if  $> 40$  kg

### Preventive

Indications: frequent headaches  $> 4x/month$ , , prolonged, severe or debilitating headaches, failure of abortive therapy, and high risk for medication overuse headaches

#### Amitriptyline

- should have a normal QT interval prior to use
- Gradually increase dose q 2 wks as tolerated to effect, up to 50 mg/day
- Patients with depression may experience worsening of depression and/or emergence of suicidal ideation

#### Cyproheptadine

- Generally does not work for  $\geq 10$  yrs old. May be associated with appetite stimulation and weight gain

#### Topiramate

- Can cause work finding difficulty and cognitive slowing, may be useful in patients with epilepsy

#### Verapamil

- May be useful in patients with hypertension. Monitor BP when initiating and with each dose increase.
- Use in patients  $> 30$ kg. Start on 40 mg tid and titrate to effect. Once on effective dose, switch to extended release formulation if available.

#### Menstrual migraine

- For female patients with migraines on oral contraceptives, a low dose (35 microgram ethinyl-estradiol or less) monophasic oral contraceptive should be used as there is increase risk in venous thromboembolism and ischemic stroke.
- The use of oral contraceptives to prevent migraine is not clearly supported in studies or in the medical literature.
- Triptans are contraindicated in patients with cardiovascular disease, uncontrolled HTN, basilar migraine, and hemiplegic migraine.

#### Non-Prescription Preventive Supplements

- Butterbur has good evidence in adult populations for headache prevention and is promising in the pediatric populations with open label studies. There is risk for hepatotoxicity if improperly prepared. Two widely used preparations that appear to be safe include Petadolex (manufacturer Weber & Weber) and Swanson Superior Herbs Butterbur Extract. May be useful in patients with allergic rhinitis/conjunctivitis. **Dosing: 75 mg oral daily**
- Riboflavin has moderate evidence in adult populations for headache prevention and is potentially effective in the pediatric population. **Dosing: 400 mg oral daily**

Opioid containing medications (e.g. Vicodin and Acetaminophen with Codeine) **should not** be used for abortive treatment of headaches

**For all medication dosing, see order sets**

# Headache Treatment by Type

## Tension Headache

**Preventive**

**Amitriptyline**

- Should have a normal QT interval prior to use
- Gradually increase dose q 2 wks as tolerated to effect, up to 50 mg/day
- Patients with depression may experience worsening of depression and/or emergence of suicidal ideation

**Cyproheptadine**

- Generally does not work for  $\geq 10$  yrs. old. May be associated with appetite stimulation and weight gain

**Topiramate**

- Can cause word finding difficulty and cognitive slowing, may be useful in patients with epilepsy

**Abortive**

**Common OTC Analgesics**

- Indicated for mild-moderate headache.
- NSAIDS tend to be the most effective. NSAIDS dosed at 10-15 mg/kg/dose and acetaminophen at 15-20 mg/kg/dose.
- Should not be used more than twice a week to prevent possible medication overuse headache and toxicity from the analgesic.

**Non-Prescription Preventive Supplements**

- Butterbur has good evidence in adult populations for headache prevention and is promising in the pediatric populations with open label studies. There is risk for hepatotoxicity if improperly prepared. Two widely used preparations that appear to be safe include Petadolex (manufacturer Weber & Weber) and Swanson Superior Herbs Butterbur Extract. May be useful in patients with allergic rhinitis/conjunctivitis.
- Riboflavin has moderate evidence in adult populations for headache prevention and is potentially effective in the pediatric population.

## Cluster Headache

**Preventive**

**Verapamil**

- May be useful in patients with hypertension. Monitor BP when initiating and with each dose increase.
- Use in patients  $> 30$ kg. Start on 40 mg tid and titrate to effect. Once on effective dose, switch to extended release formulation if available.

**Abortive**

**Sumatriptans**

- Indicated for moderate-severe headache. Triptans are **contraindicated** in patients with cardiovascular disease, uncontrolled HTN, basilar migraine, and hemiplegic migraine.
- SUMatriptan single dose for 6-11 years old is 25 mg. Single dose for  $\geq 12$  years old is 50 mg; can increase to 100 mg if needed. Max dose is 200 mg/24 hours.
- Typically third party payers require SUMatriptan prior to using any other triptans
- Nasal SUMatriptan can be used if patient has extreme nausea at onset of headache and inability to tolerate PO. Also useful if not responding well to SUMatriptan tablets. Dosing guideline 6-12 years old 5-10 mg/dose;  $>12$  years old 10-20 mg/dose. No repeat dose.

**Other Triptans**

- Rizatriptan dosing is 5 mg  $< 40$ kg and 10 mg if  $> 40$  kg

## Analgesic Medication Overuse

- Enforce/reinforce headache hygiene
- Systematically wean off abortive medications. Prepare to feel worse in a couple of weeks
- Reinforce non-medical interventions
- Keep a headache diary
- **Continue preventive therapy** if indicated as in Tension Headache Algorithm

## Sleep Apnea Related Headache

- Polysomnography
- Refer to Pulmonary

## Post Concussive Headache

- OTC Analgesics
- Non-prescription preventive supplements
- For isolated, persistent post-concussive headaches for more than 3 months, an MRI of the brain without contrast and with SWI sequence is the preferred neuroimaging study (prefer MRI obtained at CHOC Children's)
- Consider Neurology referral

## **References**

### **Outpatient Headache Care Guideline**

Holland S, Silberstein SD, Freitag F, et al. Evidence-based guideline update: NSAIDs and other complementary treatments for episodic migraine prevention in adults: report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology*. 2012;78(17):1346-53.

Lewis DW, Ashwal S, Dahl G, et al. Practice parameter: evaluation of children and adolescents with recurrent headaches: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2002;59(4):490-8.

Lewis D, Ashwal S, Hershey A, et al. Practice parameter: pharmacological treatment of migraine headache in children and adolescents: report of the American Academy of Neurology Quality Standards Subcommittee and the Practice Committee of the Child Neurology Society. *Neurology*. 2004;63(12):2215-24.

Patniyot IR, Gelfand AA. Acute Treatment Therapies for Pediatric Migraine: A Qualitative Systematic Review. *Headache* Jan 2016, 56:49-70.

Silberstein SD, Holland S, Freitag F, et al. Evidence-based guideline update: pharmacologic treatment for episodic migraine prevention in adults: report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. *Neurology*. 2012;78(17):1337-45.